

Clinical Features And Associated Comorbidities In 200 Cases Of COVID-19 In Peshawar, Pakistan

Ihteshamul Haq¹, Mohsina Haq², Momina Haq³, Seher Obaid⁴, Susan Kakakhel⁴, Mohammad Abbas⁵, Abbas Saleem khan⁶, Naumana Rhman⁷, Ghazala Zarin Afridi⁸, Fazli Zahir⁹, Fatima Syed¹⁰, Naveed Iqbal⁹, Shahzad Shoukat¹¹, Faheem Anwar¹

¹Department Of Biotechnology and Genetic Engineering Hazara University Mansehra Kp Pakistan.

²Department of Pathology Peshawar Medical college, Peshawar and Riphah international University Islamabad.

³Department of Physiology Peshawar Medical College Peshawar and Riphah international university Islamabad.

⁴Department of Physiology Northwest School of Medicine Peshawar.

⁵Department of Medicine PRIME hospital, Peshawar medical College Warsak road Peshawar.

⁶Department Of Oral pathology Peshawar Dental college, Riphah international university, Warsak road Peshawar.

⁷Department of Pathology Khyber Medical College Peshawar Kp Pakistan.

⁸Department Of Pathology Khyber Girls medical college Hayatabad Peshawar kp Pakistan.

⁹Department of Allied Health Science Iqra National University Peshawar kp Pakistan,

¹⁰Institute Of Biotechnology and Genetic Engineering agriculture University Peshawar kp Pakistan.

¹¹Department of Immunology postgraduate school universitas airlangga surabaya 60285, Indonesia.

ABSTRACT

SARS-CoV-2 is an emerging pathogen first reported in China and rapidly spreads throughout the world. The current study focused on the prevalence of symptoms, leukocytosis, leukocytopenia, thrombocytopenia, and comorbidity in patients resulted positive for COVID-19. About 200 patients were enrolled in the study first screened by ICT and then confirmed by RT-PCR. All the patients have symptoms fever, sore throat and dry cough. However 180 (90%) patients experienced tiredness, aches and pain all over the body in 193 (96%), diarrhea in 160 (80%), conjunctivitis in 64 (32%), headache in 177 (88%), loss of taste and smell in 49 (24%), a rash on skin, or discoloration of fingers or toes in 54 (27%), shortness of breath in 12 (8%), chest pain or pressure in 16 (8%), loss of speech and movement in 4 (2%) patients. 78 (39%) patients were found having co-infections including 23 (29.5%) patients have liver infection, 14 (17.9%) were kidney patients, 16 (20.5%) were heart patients and 25 (32%) were diabetic patients. Leukocytosis were found in 76 (97%) out 78 comorbid patients, while in only 2 (2.6%) non-comorbid patients had leukocytosis. Leukopenia were detected in 122 (61%) non-morbid patients out of 200 COVID-19 patients. Thrombocytopenia were detected in 118 (59%) COVID-19 patients. The patients with previous cardiovascular diseases and other comorbid conditions may face greater risk of developing the disease into severe form. 51-60 years of individuals are at high risk of getting infection. The hematologic changes are associated with COVID-19 includes thrombocytopenia, leukocytosis, and leukocytopenia.

Keywords: SARS-COV-2; Clinical features; Comorbidities; COVID-19; RT-PCR; Leukocytosis.

1. INTRODUCTION

Corona virus infection 2019 (COVID-19) is caused by Sever Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2). Most of the coronaviruses are host specific [1]. The rapid transmission rate (R_0 value) became a serious threat to human population. The world experienced the two serious outbreaks of SARS [2] and MERS [3] in 2003 and 2012, respectively. In December 2019, an outbreak arose in china and rapidly spread throughout the world. A cohort study on 36 children revealed the common symptoms including Fever (6%), dry cough (19%) and high elevated temperature of the body. Moreover, 31% patients have decreased lymphocytes while 19% have leucopenia [4]. Leukopenia, eosinophil cytopenia and lymphocytopenia are more common in patients with COVID-19 than non COVID-19 patients [5]. COVID-19 positive patients showed relatively lower absolute white blood cell (WBC), lymphocyte and eosinophil count. However, leukopenia was developed in 2 patients out of 10 and lymphocytopenia were also reported in 2 patients [5]. A 27-year-old lady in Wuhan china with history of fever and cough showed leucopenia while later she was detected positive for COVID-19[6,7]. Environmental pollution, smoking and comorbid condition (Cardio-respiratory illness, hypertension, and diabetes mellitus) likely increase the fatality rate of COVID-19 [8,9]. COVID-19 can result in Leukopenia, thrombocytopenia, myocarditis and pneumonia. Globally the COVID-19 fatality rates are 5.6 to 15.2% [10], with a greater risk of mortality in old population and those having comorbid condition like hypertension and diabetes mellitus [11]. An analysis of 46,248 cases revealed the most prevalent comorbidities that is 17 % hypertension, 8% diabetes mellitus, 5 % cardiovascular disease and 2 % respiratory morbidity [12]. A total of 6 studies involving 1527 patients revealed the percentages of hypertension, cardiovascular disease and diabetes mellitus with COVID-19 were 17.1%, 16.4% and 9.7%, respectively. It was also observed that the incidence of cardiovascular diseases diabetes and hypertension were about 3 folds, 2 folds and 2 folds respectively, higher in severe cases than non-severe cases [13]. Liver damage with COVID-19 might be directly cause by viral infection of liver cells. Patients with 2-10% diarrhorrae and detection of viral RNA in stools and blood

samples [14] implicates the possibility of viral exposure in liver. This seems the patients with COVID-19 to have higher rates of liver dysfunction [15]. A study revealed hepatitis B infection in 23 (2.1%) patients with COVID-19, Abnormal LFT and elevated aminotransferase, alanine aminotransferase and bilirubin were also noted [16]. Existing data suggest that prevalence of liver injury is more in severe cases than mild cases of COVID-19 [17]. The current study aims to address the prevalence of symptoms, leukocytosis, leukocytopenia and thrombocytopenia and comorbidity in patients positive for COVID-19.

2. Methods and Materials

The current study was performed at Hayatabad Medical Complex, Peshawar, Pakistan and in Department of Biotechnology and Genetics Engineering Hazara University Mansehra, Pakistan. Ethical approval for the study was issued by Hazara University Mansehra Pakistan.

2.1 Enrollment of patients

A total of 200 suspected patients were enrolled in the study based on the onset of sign and symptoms related to COVID-19. Every suspected patient was asked for the symptoms before testing for COVID-19. Moreover, the comorbidities were also addressed by examining the patient's history. The patients were divided in categories with respect to their age to ease in finding the comorbidities.

2.2 ICT Assay for COVID-19

Total 5ml blood was taken from everyone in two separate EDTA tubes and was stored for further analysis. To isolate serum from blood samples, 2ml blood was transferred to Eppendorf tubes and was centrifuged at 2000rpm for 2min. The supernatant was transferred through micro-pipette in a separate tube and stored for further analysis.

The COVID rapid test was performed using serum. One drop of serum (approximately 30µl) was transferred on the test strips and one drop of buffer (approximately 40µl) was added. After few minutes colored line was appeared and results were recorded.

Appearance of two distinct red lines, one line in the control region (C) and second in test region (T), showed positive results. Only one red line appearance in the control region (C) not in test region indicated negative results. The red color concentration in the test line region (T) varies depending on the conception of Covid-19 antibodies in the specimen. Therefore, any shade of red in the test region (T) was considered positive.

2.3 RT-PCR for the Detection and Confirmation of SARS-CoV-2

Throat swab sample were taken from patients screened positive for COVID-19 by ICT strip for the extraction of viral RNA. Soon after collection the throat swab were kept in 150 µL of viral preserving solution in a collection tube. Respiratory sample RNA isolation kit (Zhongzhi, Wuhan, China) were used for the extraction of viral RNA from throat swab and this process were done within two hours of sample collection. Cell lysate of 40 µL were immediately transferred to collection tube and start vortexing for 10

sec. The collection tube is leftover for 10 min at room temperature and centrifuged at 1000 rpm for 5 min. Suspension were obtained and used for RT-PCR to detect the viral RNA of SARS-CoV-2. As shown in

figure 1 the two target genes ORF1ab and nucleocapsid protein (N), were amplified and then tested at the same time in the RT-PCR process.

Target 1 (ORF1ab):

Forward primer CCCTGTGGGTTTTACACTTAA

Reverse primer ACGATTGTGCATCAGCTGA

Probe 5'-VIC-CCGTCTGCGGTATGTGGAAAGGTTATGG-BHQ1-3'.

Target 2 (N):

Forward primer GGGGAACTTCTCCTGCTAGAAT

Reverse primer CAGACATTTTGCTCTCAAGCTG

Probe 5'-FAM- TTGCTGCTGCTTGACAGATT-TAMRA-3'.

Figure 1: Primer Sequence for ORF1ab and (N)Protien for Detection by RT-PCR

SARS-CoV-2 nucleic acid detection kit was used in RT-PCR according to the company protocol (Shanghai bio-germ Medical Technology Co Ltd). The reaction mixture contains 4 µL solution of enzyme, 12 µL buffer, 3 µL of diethyl pyrocarbonate–treated water, 4 µL solution of primer Probe and RNA template of 2 µL.

The conditions which is maintained for the assay of RT-PCR are the following; 50 °C for 15 min and 95 °C for 5 min were maintained at incubation step, denaturation were performed at 94 °C for 15 sec, 55 °C for 45 sec were the conditions for extension step. Fluorescence signal were collected at the same conditions that were maintained for extension.

Ct-value or cycle threshold value of less than 37 is considered positive while 40 or more Ct-value was defined as a negative test result. The recommendations of National Institute for Viral Disease Control and Prevention China for diagnosis were strictly followed.

2.4 Complete blood count (CBC)

The CBC was determined for patients undergoing hypo-volumetric shock condition. The blood profile was determined by ALERE-380 blood cell counter machine. The main components of blood such as WBC, Platelets, and Lymphocytes values were recorded from all suspected patients.

3. Results

The enrolled 200 patients were screened for COVID-19 by ICT strip and were confirmed by the RT PCR. Age-wise distribution of the patients were found positive for COVID-19 by ICT strip and confirmed by RT-PCR are; 12 (6%), 21 (10.5%), 33 (16.5%), 43 (21.5%), 80 (40%) and 11 (5.5%) in 1-20, 21-30, 31-40, 41-50, 51-60 and 61-70 years of age respectively as shown in figure 2.




Figure 2: Age-wise distribution of the patients found positive for COVID-19 by ICT and RT-PCR

3.1 Prevalence of Symptoms

According to the questionnaire all the patients experienced fever, sore throat and dry cough. However 180 (90%) of patients experienced tiredness, aches and pain all over the body in 193 (96%), diarrhea in 160 (80%), conjunctivitis in 64 (32%), headache in 177 (88%), loss of taste or smell in 49 (24%), a rash on skin, or discoloration of fingers or toes in 54 (27%), difficulty in breathing or shortness of breath in 12 (8%), chest pain or pressure in 16 (8%), loss of speech or movement in 4 (2%) patients as shown in figure 3. The percentages were separately find out in total for each symptom, however there are some patients who have many symptoms at a time.




Figure 3: Prevalence of Symptoms in 200 enrolled COVID-19 Patients

3.2 Prevalence of Comorbidities/ coinfections

The 200 patients were found positive for COVID-19 by RT-PCR were further examine for co-infection. 78 (39%) Patients out of 200 were found to have co-infection in which 23 (29.5%) patients have liver infection (11 (14.2%) were HBs and 12 (15.4%) were HCV positive), 14 (17.9%) were kidney patients, 16 (20.5%) were heart patients and 25 (32%) were diabetic patients.

The age-wise distribution of comorbidities along with COVID-19 are HBs; 0, 1 (1.2%), 1 (1.2%), 3 (3.8%), 5 (6.4%), 1 (1.2%), HCV; 0, 0, 2 (2.6%), 5 (6.4%), 3 (3.8%), 2 (2.6%), Kidney patients; 1 (1.2%), 2 (2.6%), 3 (3.8%), 4 (5.1%), 3 (3.8%), 1 (1.2%), heart patients; 0, 0, 0, 6 (7.7%), 7 (8.9%), 3 (3.8%), diabetic patients; 0, 0, 3 (3.8%), 6 (7.7%), 11 (14.1%) and 5 (6.4%) in 1-20, 21-30, 31-40, 41-50, 51-60 and 61-70 years of age respectively, as shown in Table No 1

Table 1: Age-wise Prevalence of Comorbidities/ coinfections

Age Years	Number of Patients	Total coinfection	HBs	HCV	Kidney patients	Heart patients	Diabetic patient
1-20	12, 6%	2, 1%	0%	0%	1, 1.2%	0%	0%
21-30	21, 10.5%	6, 3%	1, 1.2%	0%	2, 2.6%	0%	0%
31-40	33, 16.5%	10, 5%	1, 1.2%	2, 2.6%	3, 3.8%	0%	3, 3.8%
41-50	43, 21.5%	18, 9%	3, 3.8%	5, 6.4%	4, 5.1%	6, 7.7%	6, 7.7%
51-60	80, 40%	30, 15%	5, 6.4%	3, 3.8%	3, 3.8%	7, 8.9%	11, 14.1%
61 +	11, 5.5%	12, 6%	1, 1.2%	2, 2.6%	1, 1.2%	3, 3.8%	5, 6.4%
Total	200	78 (39%)	11 (14.2%)	12 (15.4%)	14 (17.9%)	16 (20.5%)	25 (32%)

3.3 Prevalence of Leukocytosis, Leukopenia and Thrombocytopenia

Leukocytosis were found in 76 (97%) out 78 comorbid patients, while in only 2 (2.6%) non comorbid patients have leukocytosis. Leukopenia were detected in 122 (61%) non morbid patients out of 200 COVID-19 patients. Thrombocytopenia were detected in 118 (59%) COVID-19 patients. Shown in **Figure 4**.




Figure 4: Prevalence of Leukocytosis, Leukopenia and Thrombocytopenia

4. Discussion

200 patients were enrolled in the study and screened for COVID-19 and then confirmed by RT-PCR. As the percentage of COVID-19 patients were found high 80% in the age 51-60 this is due to the weak immune system and more prevalence of comorbidity in this group. Globally the COVID-19 fatality rates are 5.6 to 15.2% [10], with a greater risk of mortality in old population and those having comorbid condition like hypertension and diabetes mellitus [11]. The lowest number of cases were detected in age group 61+ which 5.5%. This is because the patients low number in this group. Most of the patients developed symptoms of fever, throat, dry cough, shortness in breath, headache etc. the percentages are fever, sore throat and dry cough 100%, 90% of patients experienced tiredness, aches and pain all over the body in 96%, diarrhea in 80%, conjunctivitis in 32%, headache in 88%, loss of taste or smell in 24%, a rash on skin, or discoloration of fingers or toes in 27%, difficulty breathing or shortness of breath in 8%, chest pain or pressure in 8%, loss of speech or movement in 2% patients. Same findings were also observed by [4, 18]. The percentages were separately finding out in total for each symptom, however there are some patients who have many symptoms at a time.

39% Patients out of total were found to have co-infection in which 29.5% patients have liver infection (14.2% HBs and 15.4% were HCV positive). Liver abnormalities in COVID-19 patients might be due to the viral infection in hepatocytes or due to the drug toxicity and systematic inflammation [17]. Liver injuries and morbidities is more common in severe COVID patients than patients that have mild condition with the disease [15]. However, other cause of liver morbidities such as non-alcoholic fatty liver disease, autoimmune hepatitis, alcoholic intolerance, liver disease related to consumption of alcohol, and their

effect on prognosis of COVID-19 requires further justification and evaluation. Transplantation of liver also involves the risk of viral transmission from donor to recipient, this cause is already identified in the previous outbreak of SARS therefore donor testing and screening is must [18]. Our results also showed that 17.9% patients have kidney morbidities, 20.5% were heart patients and 32% were diabetic patients. Similar results were observed by [19, 20,21]. The highest 14.1% comorbidities were found in the age range 51-60. The reason is same elder people are more susceptible to diseases or it might be due to the high number patients in this age group. Hematologic changes are associated with every disease Leukocytosis were found in 97% patients out of 78 comorbid patients, while in only 2.6% non-comorbid patients have leukocytosis. Leukopenia were detected in 61% non-morbid patients out of 200 COVID-19 patients. Thrombocytopenia were detected in 59% COVID-19 patients. Similar results were reported by [5, 13]. The commodities and other associated risk must be managed as they worsen the COVID conditions and also effect the prognosis of COVID-19.

Conclusion

Fever, sore throat, dry cough and shortness of breath are more prevalent symptoms of COVID-19. The patients with previous cardiovascular diseases and other comorbid conditions may face greater risk of developing the disease into severe form. The comorbidities also effect the prognosis of COVID-19. 51-60 years of individuals are at high risk of getting infection. The hematologic changes are associated with COVID-19 which includes thrombocytopenia, leukocytosis, and leukocytopenia. Moreover, there is also a significant drop in number of WBC in COVID-19 infection. Serious steps to be taken regarding public health management and security to prevent from SARS-CoV-2 infection.

Acknowledgment

We are grateful to Hayatabad Medical Complex Peshawar, and Department of Biotechnology and Genetics Engineering Hazara university Mansehra Kp, Pakistan for providing research facilities.

Funding Source

None

Conflict of Interest

No conflict of interest

Ethical Approval

Ethical approval was approved from Ethical and Research Committee of Hazara university Pakistan.

References

1. Baud D, Qi X, Nielsen-Saines K, Musso D, Pomar L, Favre G (2020) Real estimates of mortality following COVID-19 infection. *Lancet Infect Dis*. [https://doi.org/10.1016/s1473-3099\(20\)30195-x](https://doi.org/10.1016/s1473-3099(20)30195-x)
2. CA Donnelly, AC Ghani, GM Leung, et al. Epidemiological determinants of spread of causal agent of severe acute respiratory syndrome in Hong Kong *Lancet*, 361 (2003), pp. 1761-1766

3. Fang L, Karakiulakis G, Roth M (2020) Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? *Lancet Respir Med.* [https://doi.org/10.1016/s2213-2600\(20\)30116-8](https://doi.org/10.1016/s2213-2600(20)30116-8)
4. Guan W, Ni Z, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020; (published online Feb 28.) DOI:10.1056/NEJMoa2002032
5. Guan W, Ni Z, Hu Y. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* 2020 doi: 10.1056/NEJMoa2002032. published online Feb 28. [PMC free article] [PubMed] [Cross Ref] [Google Scholar]
6. Guan WJ, Ni ZY, Hu Y et al (2020) Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med.* <https://doi.org/10.1056/NEJMoa2002032>
7. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet.* 2020; 395: 497-506
8. Huang C, Wang Y, Li X et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020 [Epub ahead of print]. Cross ref, Google Scholar
9. Kumar D, Tellier R, Draker R, Levy G, Humar A. Severe Acute Respiratory Syndrome (SARS) in a liver transplant recipient and guidelines for donor SARS screening. *Am J Transplant.* 2003; 3: 977-981
10. Li B, Yang J, Zhao F, Zhi L, Wang X, Liu L, Bi Z, Zhao Y. Prevalence and impact of cardiovascular metabolic diseases on COVID-19 in China. *Clinical Research in Cardiology.* 2020 Mar 11:1-8.
11. Li LQ, Huang T, Wang YQ, Wang ZP, Liang Y, Huang TB, Zhang HY, Sun W, Wang Y. COVID-19 patients' clinical characteristics, discharge rate, and fatality rate of meta-analysis. *Journal of medical virology.* 2020 Jun;92(6):577-83.
12. Qamar, Z., Anwar, F., Ahmad, R., Haq, I., Khan, A. M. K., Hussain, R., ... & Khan, J. (2021). Prevalence of Hepatitis C virus and determination of its genotypes in subjects of Tehsil Daggar District Buner, KP, Pakistan. *Clinical Epidemiology and Global Health*, 12, 100809.
13. Anwar, F., Tayyab, M., Haq, I., & Shah, O. U. (2021). Viral overload of COVID-19 pandemics: Overweight people a soft target to get an infection. *International Journal of Clinical Virology*, 5(2), 070-071.
14. Haq, I., Muhammad, A., Fazli Zahir, M. K., Anwar, F., Akhtar, M. S., & Ullah, F. (2020). Serological and Epidemiology study of *Helicobacter pylori* infection among Dyspeptic patients in District Peshawar Pakistan. *Adv. Biores*, 11(3), 81-85.
15. Rehman, A., Haq, I., Asghar, M., Afridi, G. Z., & Faisal, S. (2020). Sero-epidemiological Identification of Dengue Virus in Individuals at District Shangla, Khyber Pakhtunkhwa. Pakistan. *J Biomedical Sci*, 9(3), 10.
16. Jan, Z., Ahmad, S. U., Amara Qadus, Y. A., Sajjad, W., Rais, F., Tanveer, S., ... & Haq, I. (2021). 19. Insilico structural and functional assessment of hypothetical protein L345_13461 from *Ophiophagus hannah*. *Pure and Applied Biology (PAB)*, 10(4), 1109-1118.
17. Asif, A., Asghar, M., Khan, H. U., Haq, I., Shuaib, S. L., Khalid, F., ... & Rehman, N. (2021). Antibiotic susceptibility pattern of clinical isolates of methicillin resistant staphylococcus aureus in Peshawar, Pakistan. *Annals of the Romanian Society for Cell Biology*, 25(6), 20116-20131.
18. Anwar, F., Khan, M., Salman, M., Ahmad, S., Ullah, F., Khan, J., ... & Abbas, M. (2021). Seroprevalence of hepatitis B virus in human population of district Buner Khyber Pakhtunkhwa Pakistan. *Clinical Epidemiology and Global Health*, 10, 100688.

19. Ahmad, S. U., Khan, M. S., Jan, Z., Khan, N., Ali, A., Rehman, N., ... & Zahir, F. (2021). Genome wide association study and phylogenetic analysis of novel SARS-COV-2 virus among different countries. *Pakistan Journal of Pharmaceutical Sciences*, 34(4).
20. Khalid, F. (2021). Comparative Diagnostic Analysis and Biochemical Profile in Patients with Covid-19, Dengue and Acute Febrile Illness: Suggestions for Patient Controlling. *Annals of the Romanian Society for Cell Biology*, 25(7), 1733-1744.
21. Bashir, Z., Ahmad, S. U., Kiani, B. H., Jan, Z., Khan, N., Khan, U., ... & Mahmood, T. (2021). Immunoinformatics approaches to explore B and T cell epitope-based vaccine designing for SARS-CoV-2 Virus. *Pak. J. Pharm. Sci*, 34(1), 345-352.
22. ul Haq, I., Khan, M., Rehman, Z., Anwar, F., Ullah, H., & Ullah, N. (2018). HBV prevalence in the volunteer blood donors in Peshawar, Khyber Pakhtunkhwa Pakistan. *Int J Biosci*, 13(5), 50-54.